


FBB + Drip UV

Inner Size : 8.3 x 8.3 x 2.0 cm

gProgyl

gProgyl

Clindamycin, Clotrimazole & Tinidazole
Soft Gelatin Vaginal Capsules



7 Soft Gelatin Vaginal Capsules

Only for Vaginal Insertion.
Not for oral use.

gProgyl

<https://vnras.com/drug/>

gProgyl

Clindamycin, Clotrimazole & Tinidazole Soft Gelatin Vaginal Capsules

Each Soft Gelatin Vaginal Capsule Contains :

Clindamycin Phosphate	BP	100 mg
Eq. to Clindamycin		100 mg
Clotrimazole	BP	100 mg
Tinidazole	BP	100 mg
Excipients		q.s.

Warning : If Irritation persists or increased, discontinue the use & consult the physician.


**ONLY FOR VAGINAL INSERTION
NOT FOR ORAL USE.**

Approved Colours used in capsule shells.

Dosage & Administration : One capsule should be inserted in to the vagina for 7 consecutive nights preferable before retiring to bed.

Treatment should be time so as to avoid the menstrual period.
Please read insert before use.

Storage: Store before 30°C in a cool & dry place.
Protect from direct light, heat & moisture.
Keep out of reach of children



Barcode

Mfg. Lic. No.:

Batch No. :

Mfg. Date :

Exp Date :

Mfd. By :

RENEWED LIFE SCIENCES
WHO-GMP & ISO 9001: 2015 Certified Co.
Plot No.10 & 11, Sector 6-B, SIDCUL, I.I.E
Haridwar 249403, Uttarakhand (INDIA).

gProgyl

gProgyl

HƯỚNG DẪN SỬ DỤNG

Viên đặt âm đạo GProgl



Thành phần: Clindamycin phosphate tương đương Clindamycin 100 mg; Clotrimazol 100 mg; Tinidazol 100 mg; Tá dược vừa đủ.

Quy cách đóng gói: Hộp 7 viên nang mềm.

Thời hạn sử dụng: 36 tháng kể từ ngày sản xuất.

Công dụng:

Hỗ trợ điều trị các trường hợp bị nhiễm trùng âm đạo gây ra bởi nấm Candida, nấm âm đạo, viêm âm đạo do vi khuẩn có khí hư, nhiễm trùng hỗn hợp, nhiễm trùng Leucorrhoea.

Đối tượng sử dụng: Trẻ em từ 13 tuổi trở lên và người lớn.

Hướng dẫn sử dụng:

Cách dùng:

Rửa sạch vùng âm đạo trước khi dùng.

Lấy 1 viên đặt, đặt nhẹ nhàng vào trong âm đạo, tốt nhất trong tư thế nằm ngửa, đặt càng sâu càng tốt để tránh viên đặt bị tuột ra ngoài. Tốt nhất nên đặt trước khi đi ngủ vào ban đêm.

Liều dùng:

Đặt 1 viên/ ngày, liên tục 7 ngày vào trong âm đạo trước khi đi ngủ. Tránh thời gian có kinh nguyệt.

Không sử dụng nếu có mẫn cảm với bất kỳ thành phần nào của sản phẩm.

Đọc kỹ hướng dẫn sử dụng trước khi dùng.

Cảnh báo:

Nếu xuất hiện kích ứng, ngưng sử dụng và hỏi ý kiến bác sĩ.

Chỉ sử dụng đặt âm đạo, không được uống.

Bảo quản:

Bảo quản ở nơi thoáng mát và khô ráo, nhiệt độ không quá 30°C.

Tránh ánh sáng mặt trời, nguồn nhiệt và độ ẩm.

Đề xa tầm tay trẻ em.

Sản xuất tại: Renowed life sciences.

Địa chỉ: Plot No.10 - 11, Sector 6-B, IIE, SIDCUL, Haridwar-249403, Uttarakhand, India.

Thương nhân chịu trách nhiệm về chất lượng sản phẩm:

CÔNG TY TNHH DƯỢC PHẨM & THIẾT BỊ Y TẾ TIPHACO

Địa chỉ: Thôn Thanh Xuân, TT Gia Khánh, Huyện Bình Xuyên, Tỉnh Vĩnh Phúc, Việt Nam

<https://vnras.com/drug/>

Clindamycin, Clotrimazole & Tinidazole Soft Gelatin Vaginal Capsules

GProgyl

Each Soft Gelatin Capsule Vaginal Contains:

Clindamycin Phosphate B.P.	100 mg
Eq. to Clindamycin	
Clotrimazole B.P.	100 mg
Tinidazole B.P.	100 mg
Excipients	q.s.

Approved Colours used in capsule shell.

Properties: Clotrimazole is an imidazole and has broad spectrum anti fungal action. It brings about its therapeutic effect by inhibiting ergo sterol synthesis. Inhibition of ergo sterol synthesis alters the permeability of the cell membrane of sensitive fungi and leads to structural and functional impairment of the cytoplasmic membrane. It has fungicidal action against *Candida albicans*, other species of *Candida* as well as trichomonas action.

Clindamycin is a lincosamide antibiotic primarily acting against Gram positive and wide range of anaerobic bacteria. It is congener of lincomycin. Clindamycin inhibits bacterial protein synthesis by its action at the bacterial ribosome. The antibiotic binds preferentially to the 50s ribosomal subunits and affects the process of protein synthesis. Clindamycin has antimicrobial activity against bacteroides spp., *Gerdenrella vaginalis*, *Mobiluncus* spp., *Mycoplasma hominis*, *peptostreptococcus* spp., *Haemophilus* spp., and *Corynebacterium*.

Tinidazole is an antibiotic that is used to treat certain types of vaginal infections (bacterial vaginosis, trichomoniasis). It is also used to treat certain types of parasite infections (giardiasis, amebiasis). It works by stopping the growth of certain bacteria and parasites. This antibiotic treats only certain bacterial and parasitic infections.

Indications: Infective Leucorrhoea, mixed infection and non specific vaginitis. Vaginal candidiasis, Bacterial vaginosis, Trichomoniasis, *Cadida vulvitis* & vaginal mycoses.

Contraindication: Polvag is a contraindicated in women with a history of hypersensitivity to clotrimazole (or azole group) and Clindamycin (or lincosycin group). Polvag is also contraindicated in women with a history of regional enteritis, ulcerative colitis or "antibiotic associated" colitis (pseudo membranous colitis).

Warnings and precautions: The patient should be instructed not to engage in vaginal intercourse or use of other vaginal products (such as tampons or douches) during treatment with clindamycin.

If there is a lack of response to Polvag, appropriate microbiological studies should be repeated to confirm the diagnosis and rule out other pathogens before instituting another course of therapy.

Clindamycin is found in very low levels in the serum (5%) following multiple topical applications. No systemic accumulation is seen.

Intravaginal administration of clotrimazole in human has negligible systemic absorption (3-10%)

Thus with Polvag no systemic effects are anticipated.

Polvag is to be administered intravaginally and is not to be taken orally.

Pregnancy and Lactation: Experimental and clinical investigations given on indication of harmful effects to the mother or child when clotrimazole is used during pregnancy. However, there are no adequate and well controlled studies of use of clindamycin in pregnant women. Hence Polvag should be used during pregnancy only if it is a must (under the advice of physician).

Polvag may be used in lactation with caution.

Adverse effects: Polvag vaginal capsules are well tolerated locally i.e. intravaginally. In isolated cases, vaginal irritation, itching, burning, dyspareuria and in rare cases slight increase in urinary frequency and burning or irritation intolerance and in such cases the treatment has to be stopped. Hypersensitivity reactions may occur.

Important: Only vaginal insertion. Not for oral use.

Pregnant, Lactating women should strictly abide by the instructions of their physicians.

Presentation of sales pack: Pack of 7 soft gelatin vaginal capsules.

Application and dosage:

One capsule should be inserted into the vagina, for 7 consecutive nights preferably before retiring to bed. Treatment should be timed so as to avoid the menstrual period.

Directions for use:

1. Wash hands thoroughly before and after insertion of Polvag in the vagina.
2. The capsule should be inserted as deep into the vagina as possible. This is best achieved when lying on the back with legs pulled in a little towards the body.
3. After insertion of Polvag, no activity should be done such as standing, walking running etc. hence Polvag should be inserted while retiring to bed.
4. For best results take the complete therapy for 7 days. Do not discontinue in the middle.
5. Discontinue only if severe irritation is experienced only after consulting the doctor.
6. If a particular dose is missed, administer immediately when recalled.

Mfg Lic. No:

Mfd. by :

Renowed Life Sciences

WHO-GMP & ISO 9001:2015 Certified Co.

Plot No. 10 & 11, Sector 6-B, SIDCUL, I.I.E

Hariidwar 249403, Uttarakhand (INDIA)